

Table 20D
2002 Pregnancy Nutrition Surveillance ⁽¹⁾
Nation

Page: 28
Run Date: 12 / 31 / 03

Summary of Trends in Maternal Health Indicators by Race/Ethnicity

Summary of Trends in Maternal Health Indicators by Race/Ethnicity ⁽²⁾											
Race/Ethnicity and Year		Pregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number	%
White, Not Hispanic	2002	330,578	15.1	40.3	283,276	22.8	33.9	48,677	24.9	275,055	27.6
	2001	334,613	15.6	39.7	284,413	23.1	33.6	47,298	24.7	282,932	27.5
	2000	326,763	16.1	38.7	277,756	23.3	33.5	48,052	24.3	287,217	27.8
Black, Not Hispanic	2002	161,808	10.2	48.9	135,070	28.5	26.8	26,664	45.4	118,633	51.7
	2001	165,886	10.6	48.1	135,289	28.7	27.0	25,215	43.4	122,168	50.9
	2000	165,650	11.0	47.0	129,986	29.1	27.1	27,388	44.0	127,572	51.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 20D
2002 Pregnancy Nutrition Surveillance ⁽¹⁾
Nation

Page: 29
Run Date: 12 / 31 / 03

Summary of Trends in Maternal Health Indicators by Race/Ethnicity

Summary of Trends in Maternal Health Indicators by Race/Ethnicity ⁽²⁾											
Race/Ethnicity and Year		Pregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number	%
Hispanic	2002	132,292	8.5	40.2	111,768	29.4	28.0	19,354	25.8	98,835	36.4
	2001	126,919	8.9	39.1	103,601	28.7	28.5	17,309	24.2	94,828	36.8
	2000	113,113	9.3	38.5	88,882	29.1	28.5	17,852	23.8	89,670	36.5
American Indian/Alaskan Native	2002	7,911	8.5	53.4	5,723	26.2	28.5	1,318	28.8	6,497	35.4
	2001	7,328	8.5	51.6	5,479	27.5	27.7	1,242	25.9	6,138	34.8
	2000	6,780	9.2	53.5	4,839	28.6	25.7	1,156	28.0	5,905	38.7

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 20D
2002 Pregnancy Nutrition Surveillance ⁽¹⁾
Nation

Page: 30
Run Date: 12 / 31 / 03

Summary of Trends in Maternal Health Indicators by Race/Ethnicity

Summary of Trends in Maternal Health Indicators by Race/Ethnicity ⁽²⁾											
Race/Ethnicity and Year		Pregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number	%
Asian/Pacific Islander	2002	18,419	21.3	27.1	15,938	27.8	28.9	2,838	25.3	13,514	35.7
	2001	18,300	21.3	27.3	15,441	29.0	28.3	3,003	24.0	13,048	34.0
	2000	17,653	22.5	26.3	14,653	30.7	26.5	3,104	23.6	14,211	33.0
All Other/Unknown	2002	7,412	13.0	37.3	6,291	26.5	29.8	1,220	28.5	6,197	33.7
	2001	7,722	12.5	38.4	6,208	28.9	27.7	1,325	28.2	5,217	34.7
	2000	6,130	14.1	34.3	4,936	25.3	30.2	1,036	25.9	4,931	38.8

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 20D
2002 Pregnancy Nutrition Surveillance ⁽¹⁾
Nation

Page: 31
Run Date: 12 / 31 / 03

Summary of Trends in Maternal Health Indicators by Race/Ethnicity

Summary of Trends in Maternal Health Indicators by Race/Ethnicity ⁽²⁾											
Race/Ethnicity and Year		Pregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number	%
Total All Race/Ethnic Groups	2002	658,420	12.7	42.2	558,066	25.7	30.8	100,071	30.6	518,731	35.1
	2001	660,768	13.1	41.4	550,431	25.8	30.7	95,392	29.6	524,331	35.0
	2000	636,089	13.7	40.6	521,052	26.0	30.7	98,588	29.7	529,506	35.3

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.